

**ANTIBIOPROFILAXIA IN ORTOPEDIE**

AFECTIUNE		MEDICAMENT	POSOLOGIE	DURATA
<b>Endoprotezare(sold/genunchi)</b> Infectie cu S. Epidermidis sau S. Aureus  *fara alta patologii asociata		Ceftriaxone	1g x 2/zi	48 de ore
		Cefuroxime	1,5g (intraoperator) si 0,75g/8 ore	
		In caz de alergii, Vancomicina	1g/12 ore	
<b>Fracturi inchise-</b> infectie cu S.Epidermidis, S. Aureus, Streptococ si E.Coli  *fara alta patologii asociata		Ceftriaxona	1g/12 ore	24 de ore
		Cefuroxime	0,75g/8 ore	
		Clindamicina	0,6/8 ore	
<b>Fracturi deschise-</b> infectie cu S.Epidermidis, S. Aureus, Streptococ, E.Coli si germeni anaerobi  *fara alta patologii asociata		Amoxicilina+Acid clavulanic		7 zile
		Clindamicina+ Gentamicina		
		Vancomicina+Gentamicina		
<b>Artroscopie</b>	Meniscectomie	Fara antibioprofilaxie		
	Meniscectomie +Ligamentoplastie	Ceftriaxona	1g/12 ore	48 de ore
		Cefuroxim	1,5g intraoperator si 0,75g/8 ore	
<b>Amputatie-</b> Infectie cu S. Aureus, Clostridia sau bacilli gram negativi		Clindamicina + Gentamicina	0,6 x 3/zi  5mg/kgc/zi	